## 5 CLAIMS

10

15

20

25

30

35

## WHAT IS CLAIMED IS:

- 1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95.0% identical to a sequence selected from the group consisting of:
- (a) a polynucleotide fragment of SEQ ID NO:1 or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No: PTA-3161, which is hybridizable to SEQ ID NO1;
- (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:2 or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No: PTA-3161, which is hybridizable to SEQ ID NO:1;
- (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:2 or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No: PTA-3161, which is hybridizable to SEQ ID NO:1;
- (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:2 or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No: PTA-3161, which is hybridizable to SEQ ID NO:1;
- (c) a polynucleotide encoding a polypeptide of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: PTA-3161, which is hybridizable to SEQ ID NO:1, having potassium channel beta subunit activity;
  - (f) a polynucleotide which is a variant of SEQ ID NO:1;
  - (g) a polynucleotide which is an allelic variant of SEQ ID NO:1;
- (h) an isolated polynucleotide comprising nucleotides 124 to 1095 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 325 of SEQ ID NO:2 minus the start codon;
- (i) an isolated polynucleotide comprising nucleotides 121 to 1095 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 1 to 325 of SEQ ID NO:2 including the start codon;
- (j) a polynucleotide which represents the complimentary sequence (antisense) of SEQ ID NO:1; and
- (k) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(j), wherein said polynucleotide does not

10

15

20

25

30

- 5 hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.
  - 2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a human potassium channel beta subunit protein.
  - 3. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
    - 4. A recombinant host cell comprising the vector sequences of claim 3.
    - 5. An isolated polypeptide comprising an amino acid sequence at least 95.0% identical to a sequence selected from the group consisting of:
  - (a) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-3161;
  - (b) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-3161, having potassium channel beta subunit activity;
  - (c) a polypeptide domain of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-3161;
    - (d) a polypeptide epitope of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-3161;
    - (e) a full length protein of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-3161;
      - (f) a variant of SEQ ID NO:2;
      - (g) an allelic variant of SEQ ID NO:2;
      - (h) a species homologue of SEQ ID NO:2;
    - (i) a polypeptide comprising amino acids 2 to 325 of SEQ ID NO:2, wherein said amino acids 2 to 325 comprise a polypeptide of SEQ ID NO:2 minus the start methionine;
      - (i) a polypeptide comprising amino acids 1 to 325 of SEQ ID NO:2; and
    - (k) a polypeptide encoded by the cDNA contained in ATCC Deposit No. PTA-3161.
- 6. The isolated polypeptide of claim 5, wherein the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

10

15

25

30

Ç)

- 5 7. An isolated antibody that binds specifically to the isolated polypeptide of claim 5.
  - 8. A recombinant host cell that expresses the isolated polypeptide of claim 5.
    - 9. A method of making an isolated polypeptide comprising:
  - (a) culturing the recombinant host cell of claim 8 under conditions such that said polypeptide is expressed; and
    - (b) recovering said polypeptide.
    - 10. The polypeptide produced by claim 9.
  - 11. A method for preventing, treating, or ameliorating a medical condition, comprising the step of administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 5 or the polynucleotide of claim 1.
  - 12. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:
- (a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and
  - (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.
  - 13. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:
  - (a) determining the presence or amount of expression of the polypeptide of claim 5 in a biological sample; and
  - (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.
  - 14. An isolated nucleic acid molecule consisting of a polynucleotide having a nucleotide sequence selected from the group consisting of:
    - (a) a polynucleotide encoding a polypeptide of SEQ ID NO:2;
    - (b) an isolated polynucleotide consisting of nucleotides 124 to 1095 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 325 of SEQ ID NO:2 minus the start codon;

5

10

15

20

25

30

35

- (c) an isolated polynucleotide consisting of nucleotides 121 to 1095 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 325 of SEQ ID NO:2 including the start codon;
- (d) a polynucleotide encoding the K+betaM6 polypeptide encoded by the cDNA clone contained in ATCC Deposit No. PTA-3161; and
- (c) a polynucleotide which represents the complimentary sequence (antisense) of SEQ ID NO:41.
  - 15. The isolated nucleic acid molecule of claim 14, wherein the polynucleotide comprises a nucleotide sequence encoding a human potassium channel beta subunit protein.
  - 16. A recombinant vector comprising the isolated nucleic acid molecule of claim 15.
    - 17. A recombinant host cell comprising the recombinant vector of claim 16.
  - 18. An isolated polypeptide consisting of an amino acid sequence selected from the group consisting of:
    - (a) a polypeptide fragment of SEQ ID NO:2 having potassium channel beta subunit activity;
    - (b) a polypeptide domain of SEQ ID NO:2 having potassium channel beta subunit activity;
      - (c) a full length protein of SEQ ID NO:2;
    - (d) a polypeptide corresponding to amino acids 2 to 325 of SEQ ID NO:2, wherein said amino acids 2 to 325 comprise a polypeptide of SEQ ID NO:2 minus the start methionine;
      - (e) a polypeptide corresponding to amino acids 1 to 325 of SEQ ID NO:2; and
  - (f) a polypeptide encoded by the cDNA contained in ATCC Deposit No. PTA-3161.
  - 19. The method for preventing, treating, or ameliorating a medical condition of claim 11, wherein the medical condition is selected from the group consisting of a gastrointestinal disorder, a reproductive disorder, an immune disorder, a neural disorder, a cardiovascular disorder, a pulmonary disorder, a disorder related to hyper potassium channel activity, an immune disorder related to aberrant NF-kB

activity, pineal gland associated disorders, migraine headaches, disorders associated with aberrant melatonin synthesis and/or release, delayed sleep phase syndrome, aberrations in circadian cycle, mammary cancer tumorigenesis, disorders associated with low DNA repair capacities or low free-radical buffering capacity, sleep disorders, age related disorders associated with descreased melatonin secretion, and cancer.